

**Deciduous
(primary)
Usual age of
eruption**

**Permanent
(colored blue)
Usual age of
eruption**

- Central incisor (8–10 months)
- Lateral incisor (8–10 months)
- Canine (cuspid) (16–20 months)
- 1st molar (15–21 months)
- 2nd molar (20–24 months)

- 2nd molar (20–24 months)
- 1st molar (15–21 months)
- Canine (cuspid) (16–20 months)
- Lateral incisor (15–21 months)
- Central incisor (6–9 months)

- Central incisor (7th year)
- Lateral incisor (8th year)
- Canine (cuspid) (11th–12th year)
- 1st premolar (9th year)
- 2nd premolar (10th year)
- 1st molar (6th year)
- 2nd molar (12th–13th year)
- 3rd molars (17th–25th year)

- 2nd molar (12th–13th year)
- 1st molar (6th year)
- 2nd premolar (10th year)
- 1st premolar (9th year)
- Canine (cuspid) (11th–12th year)
- Lateral incisor (8th year)
- Central incisor (7th year)

Incisive fossa

Palatine process of maxilla

Horizontal plate of palatine bone

Greater and lesser palatine foramina

Central incisors

Lateral incisors

Canines

1st premolars

2nd premolars

1st molars

2nd molars

3rd molars

Upper permanent teeth

Lower permanent teeth

F. Netter M.D.